


Assisted Hygiene and Wellness solutions



Introduction

Since the creation of Arjo over sixty years ago, improving the hygiene care experience for patients, residents and caregivers has been at the very heart of what we do.

Our journey began when Arne Johansson, the original company founder, built the first height adjustable bathtub driven by a desire to improve care for his wife who had impaired mobility. This innovation was the first of many, all designed to improve safety and efficiency for busy caregivers, while focusing on the dignity, comfort and wellness of residents and patients during the showering and bathing process. This clear vision led to Arjo becoming the global market leader in the provision of hygiene solutions, a position it retains today.

We recognise that clinical decision making is complex and should be based upon the best available evidence. We invite you to review this clinical resource providing examples of evidence supporting the use of hygiene and bathing solutions.

Assisted hygiene and wellness

Hygiene care is more than just keeping the skin clean. For many people having the opportunity to shower or bathe can have a significant impact on their general well being and quality of life. For a patient during either a short or long term admission to a healthcare facility, continuing their personal hygiene habits often established over a lifetime can be challenging. Time pressured staff, cold unwelcoming hygiene facilities and often rigid routines means focus is frequently on getting the person clean as quickly as possible rather than on person centred hygiene care that takes account of individual preferences. Providing personalised hygiene care can lead to a more relaxing and enjoyable hygiene experience with benefits for the persons wellbeing.



Assisted hygiene care challenges

Hygiene care tasks place a significant and increasing burden on time pressured caregivers as the number of frail, elderly, heavy or dependent residents and patients with complex health issues continues to rise.¹ Decisions regarding how people access hygiene facilities are made each day, decisions regarding the level of assistance required by individual patients and the choices that are made can impact patient dignity, comfort as well as safety and both the physical load, and workload for caregivers.



Caregivers

Showering and bathing tasks expose caregivers to a high risk of back and shoulder pain and injury.² Lack of appropriate showering equipment increases the risk for inefficiency and physical strain experienced by staff. Efficiency of providing hygiene care is dependent upon the mobility level of resident or patient, type of equipment available and appropriate training received.³ Perception that taking a person to the bath/shower takes longer and is more physically demanding compared to washing the person within their own bed.



Residents and Patients

Showering or bathing can have a significant impact on a person's wellbeing, but hygiene events are often rushed with a focus on getting the patient or resident clean as quickly as possible. Risk of falls on wet bathroom floors can be particularly hazardous especially for those with restricted mobility,⁴ or already classed as at risk from falls. Risk for cross infection of using shared bathtubs without robust decontamination. For preventative skin care the need to keep the skin clean.⁵ Maintenance of dignity during the hygiene and bathing process.



Facility

The need for healthcare efficiencies and lack of storage space within facilities often means that under-utilised bathrooms are used as storage spaces for equipment, impacting the working space available to move hoists and other equipment required for the hygiene tasks. Space within bathroom areas can be small and not ergonomically designed for people with increasing demands for assistance during hygiene care. Limited space for transfer, a lack of appropriate lifting and showering equipment, and a lack of clear processes and protocols for assisted hygiene activities may increase the risks of patient and caregiver injury.

Person centered hygiene care

When entering a care environment for a short or long term stay, reliance on others to assist with personal hygiene may have a significant and potentially detrimental effect on the individual, who should always be involved in decisions regarding how their hygiene needs will be met.

Caregivers need to assess if a patient or resident can bathe or shower independently, if help is required and if so the type of assistance needed including equipment that may be required to support the activity in safety and with dignity.

Depending on the facilities available and the care needs of the individual, choices for providing hygiene care are centred on *showering, bathing in a tub or bathing within or at the bedside.*

Decisions that are made are often influenced by time available by the caregiver and the perceived ease of carrying out the different hygiene procedures. The quality of care received and the quality of work performed by the caregiver is reliant upon having the right environment, using the right equipment and utilising the right care skills. This is represented by the **Arjo Positive Eight**.

The Positive Eight

Promoting resident/patient mobility is the driving force behind the Positive Eight – our philosophy for generating long-term benefits for patients, caregivers and care facilities. Investing in the right environment, suitable assistive equipment and adopting the best caregiver practices can provide the optimal conditions to promote and enhance mobility. Promoting mobility can reduce the need for support, reduce caregiver injuries and improve efficiency. Ultimately this all leads to improved care and financial outcomes.




② Improved vital functions

① Mobility

⑤ Reduced need for support

⑥ Reduced injuries and improved efficiency



③ Reduced consequences of immobility

④ Quality of life

⑧ Improved care and financial outcomes

⑦ Reduced sick leave, turnover and compensation claims

The right **environment, equipment and care skills** need to be in place to allow the benefits of the Positive Eight to flow



Supporting care choices

Having the right hygiene equipment for residents and patients who are self-reliant will support their continued independence and help ensure well-being is maintained.

Providing assistive devices and hygiene equipment for the more dependent person will enable access to shower or bathing facilities as preferred, as well reduce the exposure to unacceptable levels of physical overload and risk of injury to the caregiver.

The Arjo Mobility™ Gallery

The Mobility Gallery, as a functional assessment tool, can be used to assist facilities and caregivers to identify the hygiene requirements of residents and patients for whom they provide care and support decisions related to showering, tub bathing or bathing at the bedside.

Different mobility levels require a range of hygiene solutions






























Arjo hygiene solutions are designed to assist in reducing risk of injury to caregivers and patients, and make hygiene tasks comfortable, efficient and dignified for the residents and patients. Our showering and bathing solutions are developed to meet both the needs of the individual, and their caregivers. Having a comprehensive range of solutions allows the caregiver and the resident/patient to have a choice to best suit their functional needs and preferences.

Assisting someone to maintain his or her personal hygiene needs can contribute to the comfort, safety, wellbeing and dignity of the individual.⁶



By matching the right hygiene solution to the needs of residents and patients, Arjo hygiene equipment can provide a level of support designed to maintain and, where possible encourage and promote mobility.

Solution chart

Mobility level	Showering	Reclining sit bath solutions	Sit bath with integrated chair	Recumbent bath solutions
 ALBERT	 Foldable seat	 Reclining sit bath	 Sit bath with integrated chair	 Bath lift chair  Recumbent bath
 BARBARA	 Foldable seat  Hi-low shower chair	 Reclining sit bath	 Sit bath with integrated chair	 Bath lift chair  Recumbent bath
 CARL	 Hi-low hygiene chair  Multi-purpose hygiene chair	 Reclining sit bath	 Sit bath with integrated chair	 Bath lift chair  Recumbent bath
 DORIS	 Multi-purpose hygiene chair	 Reclining sit bath		 Bath lift trolley  Recumbent bath
 EMMA	 Shower trolley			 Bath lift trolley  Recumbent bath

Reducing caregiver risk during assisted showering

Showering may be preferred for frequent personal hygiene procedures, and is regularly used in hospitals to prepare patients prior to surgical procedures.

Appropriate showering equipment promotes better posture and better care

Every day, caregivers assist residents and patients to take a shower. For the caregiver, the risk of physical overload from working in positions which involve bending, stooping and twisting, often using fixed height shower chairs means they can be subjected to high levels of static (postural) overload, which subsequently increases their risk of musculoskeletal injury.

The correct use of suitable adjustable height showering equipment can reduce the need for adopting awkward postures and can assist toward reducing injury related absenteeism,

as well as improve working efficiency during the showering procedure.⁷

The suitable supportive equipment will also offer a more pleasant, comfortable and dignified shower experience for the individual. 'Knibbe *et al*' undertook a laboratory study which illustrated the percentage of time caregivers spent in healthy working postures when showering patients using a variety of assistive devices.⁸ The highest percent of neutral back postures were seen when caregivers used the Carendo hygiene chair.

Percentage of observations of caregivers with neutral back postures for 6 types of equipment (N=79.968 observations, N=112 Procedures) Knibbe *et al*⁸

Equipment	% of neutral back postures (all)*	% of neutral back (mobility class C)*
High - low shower chair with profiling seat	77.3	73.9
High - low shower chair	66.4	67.1
Ergonomic high - low shower trolley	59.5	61.8
Fixed - height shower chair (normal)	66.0	57.4
Fixed - height shower chair (high)	65.9	56.4
High - low profiling bed	48.6	53.6

*Differences (Kruskal-Wallis Test) statistically significant $P < .05$,



ALBERT



BARBARA



CARL



DORIS



EMMA



Using the Carendo resulted in the caregiver adopting 77% neutral back positions compared to only 48% when washing a person within their bed.⁸

Reducing static load

A study has shown that using the Carevo, an ergonomic and adjustable height shower trolley reduces the static physical overload experienced by caregivers compared to when they use a standard shower trolley.⁹

The use of height adjustable devices when bathing or showering patients enables caregivers to work at their correct ergonomic height, reducing the static load on their musculoskeletal system and subsequent risk of injury and long term damage.

The design of the Carevo shower trolley is to enable the caregiver to get closer to the resident/patient and hence further reduce caregiver static load while showering patients. Results of these recent studies illustrate the advantages of using this equipment over washing a person under their bed and also on a standard width and static height shower trolley.





Increase efficiency in assisted showering

Using known average timings for washing in bed and washing with an appropriate shower device¹⁰ a study undertaken in a French Long Term Care facility showed that a total of 314 minutes or 5.2 hours could be saved daily if carers used appropriate height adjustable hygiene equipment instead of washing C, D and E patients within their bed. This equated to a potential annual saving of 20,000 euros

Enabling efficient working

Washing, showering and cleaning patients is one of the most time-intensive care activities that caregivers can perform. Research showed that carers in a nursing home devoted 22% of their total working hours to washing, dressing wounds, showering, bathing and


dressing.¹⁰ With a shortage of caregivers and increasing dependency levels of residents and patients, finding more efficient ways of working has never been more important.

Time saving

Although there is a common perception that washing a resident or patient whilst they are within their bed is the fastest method, a recent study has shown that **using a height adjustable shower chair, a profiling hygiene chair or a shower trolley with 'C' mobility level residents and patients can be 20% - 35% more efficient.**¹⁰

Improved efficiency

The use of appropriate hygiene equipment often enables caregivers to work more efficiently, reduce the number of transfers and potentially work without the assistance of a second caregiver. The one-to-one person focused interaction between patient/resident and caregiver is perceived as important in relation to aiding communication and reducing residents and patients stress, especially when caring for people with dementia.¹¹




Although there is a widespread belief that washing a resident or patient in bed is faster, a heightadjustable shower chair can be 35% more efficient.

Source: Schimmel G, Knibbe NE, The Smart Shower Chair, Final Report of the Zorg voor Beter (Better Care) Smart Shower Chairs project, 2005.


■ Washing in bed
■ Washing in Carino
■ Washing in Carendo
■ Washing in Carevo

When using a Carendo multipurpose hygiene chair with a C patient there is a potential of saving 14 minute for each showering event, when compared to washing a C patient within their bed¹⁰


Total number of transfers



Transfers performed with 2 caregivers



Number of transfers performed in an unhealthy way




Increased efficiency was demonstrated in an observation study

A study in Denmark¹¹ identified 64% of the transfers as potentially risky and unhealthy for the staff, of which 74% were in the hygiene area. The study pointed to one main gap related to time consumption and ergonomics, which limited access to ergonomic hygiene equipment for showering. This resulted in the organisations implementation of the Carendo hygiene chair, and together with a detailed training programme the follow-up after sixteen months showed an **83% reduction of transfers performed in an unhealthy way**. The total number of transfers had decreased by 39%, and the number of transfers requiring two members of staff was eliminated. Members of staff reported to be more satisfied since unnecessary or potentially dangerous transfers had substantially decreased, due to a more


efficient way of working. The investment in the Carendo Multipurpose hygiene chairs saved the equivalent time of one full time equivalent caregiver.

In a long term care facility 78% of caregivers agree that using Carendo improves their ability to perform washing and hygiene of all areas of the body including skin folds.¹¹

TYPICAL SHOWERING WORKFLOW WITHOUT A CARENDO



TYPICAL SHOWERING WORKFLOW USING A CARENDO





The hygiene care environment

The hygiene care environment within healthcare can vary between an en-suite facility dedicated to use for one person, or a centrally located bath or shower room which is shared. Ensuring that a shared bath or shower room is suitable for a number of individuals is important. The more flexible in design, the better the environment can be adapted for individual needs, for both assisted and independent showering or bathing. The possibility to make just small adaptations can be the difference between independence and loss of functional ability.

The Architects Guidebook (2019)¹² provides further information related to both showering and bathing environments and space requirements for different mobility levels of residents and patients using a variety of assistive hygiene devices.

[To find out more please visit arjo.com](https://arjo.com)



Table of evidence – showering

Author	Design	Key Findings
Knibbe et. al. Evaluating different methods of showering and washing patients; assessing ergonomic, time and quality aspects, Am J SPH, Vol. 6, No 2, 2016.	Laboratory study. The aim of the study was to provide a quantitate insight into postural load during hygiene tasks using video analysis and survey. 112 standardised hygiene procedures performed by 4 nurses, using six different types of equipment, and two washing techniques on acting patients with varying mobility.	Patient mobility, type of equipment, and washing technique were related significantly to hygiene care duration. The skill of the nurse, type of equipment, and mobility were linked to the postural load on the back, overall, lower satisfaction with procedures on the high-low bed. Recommendations for reducing duration and extent of static load are given but more research is needed.
Hallström K. Wiese N. Improving Quality of Care and Workflow by Optimising the Working Conditions. A Case Study from a Danish Nursing Home. Proceedings 19th Triennial Congress of the IEA, Melbourne 9-14 August 2015	Case study. The aim of this study was to establish a more efficient and safer process for managing resident transfers. The objectives were to reduce the static load, and to reduce both the number of transfers classified as potentially risky or dangerous. Four physiotherapists assessed the resident transfers for a period of twenty-four hours. Mobility level classification, NIOSH equation, and static load analyze were methods used.	The baseline assessment identified 64% of the transfers as potentially risky and unhealthy for the staff, of which 74% were in the hygiene area. The observation identified limited access to ergonomic hygiene equipment for showering. A new process was implemented incl. the use of Carevo® together with ceiling lifts and training. The follow-up after sixteen months showed an 83% reduction of transfers performed in unhealthy way. The total number of transfers was decreased by 39%, and the number of transfers requiring two members of staff was eliminated.
Knibbe N.E. et. al. How Smart is the Carevo®? Results of a Study on a New Shower Trolley Generation. The Ergonomics Open Journal, 2013, 6, 1-5.	Laboratory study. The purpose of the study was to compare the traditional height adjustable shower trolley with Carevo®, a shower trolley with a new design. The study involved 4 nurses carried out 5 shower activities, with D and E patients all postures observed were filmed and analysed using OVAKO.	Differences seen when caregiver works with Carevo®, compared with the traditional trolley: 10% improvement in time spent in a neutral safe back posture, 3% improvement in time spent in a non-flexed, non-twisted back posture, 9% improvement in time spent in a neural safe neck posture. In additional more time was in the middle part which resulted in 13% reduction in static load.

Author	Design	Key Findings
Freitag S. et. al. Frequent Bending— An Underestimated Burden in Nursing Professions, Ann. Occup. Hyg., pp. 1-11, 2011.	The aim of the study was to quantify the total duration per shift in which nurses work in a forward bending position over 20°. In addition, the influence of several factors on the occurrence of sagittal trunk inclinations in nurses was investigated. Trunk postures were recorded for nine nurses from four nursing homes and 18 nurses from seven hospitals using the CUELA measurement system. A total of 79 shifts, 27 in nursing homes and 52 in hospitals, were analysed. All measurements were supported by video recordings.	The total duration of inclinations per shift was significantly affected by the working area. Nursing home nurses worked about twice as long per shift in a forward bending position compared with hospital nurses (112 versus 63 min) and they assumed almost one-third more inclinations per shift. Another factor was the extent of personal basic care, incl. hygiene tasks, performed by the nurses. It is very likely that future preventive measures, focusing on reducing the huge amount of inclination, would reduce the physical stress in everyday nursing work substantially.
Brinkhoff A., Knibbe N, The ErgoStat Program . Pilot study of an ergonomic inter- vention to reduce static loads for caregivers, www.asse.org Professional Safety, Ergonomics, May 2003	Pilot study. An evaluation of an Ergonomic intervention to reduce static loads for caregivers. Measurement of postures before and after the program using OWAS.	After the intervention, a significant increase was observed in the percentage of time that caregivers spent working in a healthy position. The most notable improvement was seen in performing direct care tasks – demonstrated in particular with the improvement in the head position.
Schimmel G., Knibbe N.E. The Smart Shower Chair Final report of the Zorg voor Beter [Better Care] Smart Shower Chairs project, Opella and Locomotion 2003.	Pilot study. An evaluation of an Ergonomic intervention to reduce static loads for caregivers. Measurement of postures before and after the program using OWAS.	The time savings as a result of using a smart shower chair, Carendo® are 14 minutes (Mobility Class C) and 6.5 minutes (Mobility Class D) per shower, and at the same time the static load is lower when using Carendo®. The threshold for working with Carendo® appears to be fairly high, according to this project. Training, instruction and working regularly with the smart shower chair is therefore necessary in order to use the aid in an optimal manner.

Reducing caregiver risk during assisted bathing



Bathing promotes health benefits and can play an important role in nurturing wellness and improving quality of life. It has been recognised as a pleasurable relaxing experience for centuries. Bathing is a basis for hygiene, in addition it can promote a sense of well-being. A warm bath can relax both muscles and mind.¹³

Bathing is becoming even more important for individuals with impaired mobility and cognitive function. During the bathing process the resident/patient can be encouraged to take part and wash as much of himself as possible. This provides an opportunity to help maintain dignity and a level of independence.

"But what are the bathing challenges for caregivers and residents?"



FOR RESIDENTS

GETTING IN AND OUT OF THE BATH IS A CHALLENGE

- The resident may feel cold if left wet and exposed during transfers
- The resident may be fearful of slipping
- Getting in and out of the bath may trigger undesirable behavior
- Dignity aspects related to undressing and dressing, this should happen in the bathroom to ensure dignity



FOR CAREGIVERS

THE BARRIER IS TIME AND SPACE

- There may be a risk of static overloaded for a caregiver
- There may be a perception that bathing takes long time
- Assisted bathing involves a high number of assisted transfers
- Often no space to dry and dress
- The facility does not have the right equipment to support or transfer the resident/patient to/from the bath or provide sufficient support whilst immersed in the water



ALBERT



BARBARA



CARL



DORIS



EMMA



Risk of caregiver injury during bathing procedures - appropriate bathing equipment promotes better posture and better care

Lifting residents and patients into and out of a bathtub, leaning over into the tub to reach the person to provide hygiene assistance as well as managing patient falls within the bathroom have long been accepted as potential causes of physical strain and cause of injury to caregivers. Choosing the right bathing equipment for the resident/patient as well as giving consideration for the needs of the caregiver will assist the organisation to pro-actively tackle the problem associated with such heavy handling and adoption of poor working postures. The extent of the physical stress experienced by the caregiver is dependent upon a number of factors, the mobility level of the resident/patient, the skill of the caregiver, the environment in which the bathing is taking place and the equipment which is available to use.^{7,14}

Enhance mobility

Maintaining and promoting mobility is a crucial part of good quality care. Immersion in warm water may ease muscle tension and increase blood circulation.^{13,15,16} Importantly, being immersed in water makes the body feel lighter, it can therefore reduce the load on joints, and movement may become easier. Bathing may offer a natural and pleasurable way to help patients and residents maintain their existing mobility as part of a mobility maintenance program. In additional bathing in warm water could be an important benefit for people with limited joint mobility and trying to regain mobility during rehabilitation.¹⁷

Bathing a resident or patient using a high low bathtub can reduce static overload to the caregivers back by over 15% when compared to using a fixed height bathtub.¹⁸

Equipment	Percentage of time caregiver spent in healthy posture. (Higher score is less harmful)
High - low chair	58,8%
High - low bath	56,6%
High - low shower trolley	53,0%
Fixed bath	41,6%
High - low bed	39,4%
Fixed shower trolley	38,6%
Fixed shower chair	31,4%
Conclusion	

With respect to postural load, the High low bath and other height adjustable equipment place significantly less stress on the musculoskeletal system, especially when compared to the fixed bath, fixed shower trolley, fixed shower chair and hi-low bed.¹⁸



Increase efficiency in assisted bathing

Time savings and bathing

Although there is very little reliable research on the time taken to assist a resident or patient to take a bath, 'time' or 'perceived lack of it' is the most frequently heard reason for not assisting residents to bathe in a tub. The whole bathing process can be divided into 3 different 'steps':

- **Pre-bathing (preparation of the environment and patient)**
- **Washing in the bathtub**
- **Post-bathing (drying and dressing)**

Knibbe et al (2016) showed in their study, that it is the caregiver that often determines the time taken for washing, depending upon the perception of the subjective feelings of the resident/patient and went on to suggest that when bathing and showering, caregivers will often make the most of the 'wellness opportunity' rather than just performing the assisted cleaning.⁸ Having functions on the bathtub which provide caregivers with time savings on the pre-bathing stage, such as pre-fill function on the tub can only assist to provide more time for positive 'well being' opportunities with their residents and patients for them whilst they are in the tub itself.

Plan for the whole bath session:

When planning the bathing area and the bathing process, it's important to keep the whole bathing session in mind. The workflow needs to be planned; from undressing, to bathing, showering, toileting and re-dressing. In the planning of a central bathing area and the bathing process several aspects need to be taken into consideration.


Plan how resident/patient gets to the bathroom

It is a good idea to be able to get to the bathroom dressed, ambulating or sitting in a wheel-chair/shower chair or bath chair. As this preserves the resident/patient dignity and ensures that the person stays warm

Where to get undressed & dressed

The preferred solution is to be undressed and dressed either in the bedroom or in the bathroom. Many residents and patients need some level of assistance; here an active lifter, a bath lift chair or trolley could be good options. Enough space, dignity and risk for the undressed person getting cold are key aspects to be kept in mind.

3 STEP BATHING WORKFLOW



Have a secluded place for getting to the toilet

It is a good idea to empty their bladder and bowel before bathing or showering. The toilet area could be integrated with, or be adjacent to, the bathroom.

Plan a safe way to enter the bathtub

Entering the bathtub should be carried out in a way that is safe for both the resident/patient and the caregiver reducing the risk of slipping and subsequent fall.

Create a safe working environment

During bathing the caregiver should be able to provide assistance to the resident/patient using in a safe working posture, the tub and patient seat should be at a comfortable height while the resident/patient relaxes comfortably.

Offer a shower

The resident/patient may prefer a shower prior to, or after the bathing session. To support independent or assisted showering as an alternative to bathing, there should be a separate showering area, this area could also be used to clean the resident/patient if in the event of the resident/patient needing to empty their bladder/bowels during the bathing session.

Getting out of the bath

Even if the resident/patient did not need support to enter the bath, exiting could be difficult. After the bathtub the resident/patient is warm, wet and maybe feeling very relaxed. Using a lifter to reduce the risk for the resident/patient falling and to avoid strain on the staff is to be recommended in most cases.

Drying and dressing

A heated area for assisted drying is good to plan for. It is important that the resident/patient can relax and doesn't get cold after the bath. A warm towel, is a welcome detail enhancing the feeling of well being and maintains the persons dignity.

Relaxation

It is also important to plan for an area where the resident/patient can sit and relax for a while. A bathing session can be nice, but also exhausting, and a short time of recovery after the bath is often appreciated.

Cleaning

To avoid the risk for cross-infection, the bathtub and equipment used during bathing or showering will need to be cleaned and disinfected. A bath or shower-panel with built-in cleaning disinfection is a welcome time-saving feature for the caregivers.

Bathing to enhance wellbeing



Reduce Pain and Induce Relaxation

Regular bathing can reduce pain when part of a pain management programme^{15,19} and also increase relaxation.^{20,21} Bathing with bathing oil can also be a gentle and effective way to clean a resident with sensitive skin.

Bathing in warm water in general has a naturally calming effect. The warm bath water supports the body in a near weightless state, helps relieve muscle tension and induces relaxation. The relaxing nature of bathing in warm water has been shown to achieve positive clinical results, potentially diminishing the need for painkillers and sedatives.²² It may also be beneficial in reducing stress and anxiety and to improve sleep.²¹

Goto et al in their randomised controlled trial suggests that routine immersion bathing has positive effects on both mental and physical health, such as lowered stress and anxiety levels as well as increasing hyperthermia action which increases blood flow.¹²



A study using the Parker® Spa bath within a palliative care environment concluded that using a spa bath with such patients was a simple way for nursing staff to enhance the comfort of their patients within the normal course of their duties and without the need for additional training.²³



The action of the bubbles bursting against the skin remove dirt particles from the skin in a gentle brushing action.^{28, 29, 30}

Stimulate the senses

Snoezelen™ multi-sensory environments are often specially designed to deliver stimuli to various senses of the person, using lighting effects, color, sounds, music, scents, etc., they are by nature, relaxing spaces that help to reduce agitation and anxiety, but they can also engage and delight the user, stimulating reactions and encouraging communication.

The bathing session could give an excellent chance for sensory stimulation, the temperature and tactile feeling from the water, the smell of bath oils, colour, light and music could be used to stimulate the resident.

The use of light, sound and smell can be used to gently stimulate the senses to create either a relaxing or stimulating environment. Music during bathing may calm the resident and lessen aggressive and agitated behaviors.²⁴ But we need to keep in mind all residents and patients are different and what works well for someone would cause anger or frustration for someone else. This means the bathing session and bath environment need to have a person focused care approach.

Using SOUND & VISION may engage the senses and potentially enhance the bathing experience for the resident/patient. It has been shown that the use of music can contribute towards the lessening of aggressive and agitated behaviors in older people with Dementia.^{24, 25}

Support skin care

The skin protects the body and bathing may help to protect the skin. However, what you put in the bathwater can complement or counteract good skin management. Making good choices in bath liquids and choosing Hydrosound® as a washing method are two factors that can contribute to optimum skin care.

Hydromassage

It is possible that HYDROMASSAGE can assist in pain management,^{26,27} in particular chronic pain, as well as contribute to the reduction of lower limb stiffness.²⁸ Hydromassage works by creating millions of small bubbles and has the effect of creating agitation in the bathwater. This in turn provides a gentle massaging effect which is intended to support relaxation as well as give a pleasant sensory experience.

Hydrosound

The use of HYDROSOUND, a low frequency ultrasound, has been shown to be able to cleanse the skin through the gentle mechanical action of streaming microbubbles. Low frequency ultrasound has also been shown to improve peripheral circulation and has been used in wound care to remove non-viable tissue.

The action of the bubbles bursting against the skin remove dirt particles from the skin in a gentle brushing action.^{30,31,32}

'adding music-assisted bathing into the activities of daily living routine may assist in lessening of aggressive and agitated behaviours exhibited prior to and during bathing'.³²

Table of evidence – bathing

Author	Design	Key Findings
Goto et. al. Physical and Mental Effects of Bathing: A Randomized Intervention Study, Evidence-Based Complementary and Alternative Medicine, Volume 2018, Article ID 9521086	<i>Randomized controlled trial.</i> This study compared the effects on health of immersion bathing, whole-body immersion bathing, and shower bathing in 38 participants who received 2-week intervention of immersion bathing in warm water (40°C) for 10 min followed by 2-week shower bathing without immersion or vice versa (n = 19 each group).	Visual analog scale scores were significantly better for fatigue, stress, pain, and smile and tended to be better for self-reported health and skin condition after bathing intervention than after showering intervention. The Health Survey showed significantly better general health, mental health, role emotional, and social functioning scores. Scores were lower for stress, tension-anxiety, anger-hostility, and depression-dejection.
Karagulle M, Real-life effectiveness of spa therapy in rheumatic and musculoskeletal diseases: a retrospective study of 819 patients Int J Biometeorol 2017;61	<i>Retrospective observational study.</i> The objective was to determine the use and efficacy of spa therapy in 819 patients with a wide spectrum of rheumatic and musculoskeletal diseases under real-life clinical practice circumstances. All adult patients with rheumatic and musculoskeletal diseases, who were prescribed a spa therapy in various health resorts in Turkey between 2002 and 2012 were analyzed. Patients followed a usual 2-week course of spa therapy. The patients were examined before and after the spa therapy.	The patients included had the following diagnoses: 536 osteoarthritis; 115 fibromyalgia; 50 lumbar disc herniation; 34 cervical disc herniation; 23 nonspecific low back pain; 22 ankylosing spondylitis; 16 rheumatoid arthritis; 9 rotator cuff tendinitis; and 14 other conditions/diseases. The result showed statistically significant decrease in pain scores in all patients except for in the subgroups hip osteoarthritis and rheumatoid arthritis. In addition the result showed a statistically significant improvement in function in all patients except for the patients suffering hip osteoarthritis, rheumatoid arthritis and rotator cuff tendinitis. The study results suggest that real-life spa therapy may be effective in a variety of rheumatic and musculoskeletal diseases by improving pain and function.
Goris D. E. et. al, Quantitative systematic review of the effects of non-pharmacological interventions on reducing apathy in persons with dementia, Journal of Advanced Nursing, Nov 2016; 72 (11)	<i>Quantitative systematic review.</i> The aim of the study was to review the quantitative evidence concerning the effects of non-pharmacological interventions on reducing apathy in persons with dementia. The CINAHL, PubMed, PSYCH info and Cochrane Trials databases were searched for research published in English, inclusive through December 2014, with no early year limiters set.	In the search, 16 international randomized controlled trials or quasi-experimental studies were identified. The review suggest that selected non-pharmacological interventions are effective in reducing apathy among persons with Alzheimer Dementia Specifically. Music therapy, tailored personal contact, cognitive stimulation therapy, multi-sensory behavior therapy (including Snoezelen), group art therapy and therapeutic conversation all show some effect in reducing apathy without producing negative side effects. Overall, these interventions are heterogeneous in many ways, including their method of engaging a person with dementia, the apparent target of their therapeutic effect (behavioral, emotional or cognitive domains of apathy) and the extent to which interventions were tailored to a specific individual.
Matzer et. al. Stress-Relieving Effects of Short-Term Balneotherapy – a Randomized Controlled Pilot Study in Healthy Adults, Forschende Komplementärmedizin 2014;21:105-110	<i>Randomized study.</i> The aim of the study was to evaluate the stress-relieving effects of balneotherapy compared to Progressive Muscle Relaxation (PMR) 49 healthy volunteers were randomized to balneotherapy, PMR, or a resting control group, each intervention lasting for 25 min; subjective relaxation and salivary cortisol were measured. It was also examined whether participants with a high versus low stress level would have a different response.	In a pre-post comparison, salivary cortisol decreased and subjective relaxation ratings increased in all 3 groups. Study participants in the balneotherapy group rated themselves as more relaxed after the intervention as compared to the other two groups. Participants with a high versus low stress showed a similar relaxation response. Findings suggest that compared to PMR and resting, balneotherapy seems to be more beneficial with regard to subjective relaxation effects and similarly beneficial with regard to a decrease in salivary cortisol.

Author	Design	Key Findings
Falagas M. E. The therapeutic effect of balneotherapy: evaluation of the evidence from randomised controlled trials, <i>Int J Clin Pract.</i> 2009 Jul;63(7)	<i>Literature Review.</i> Systematic review with the aim to evaluate balneotherapy. PubMed, Scopus and Cochrane library were searched for randomised controlled trials, examining the clinical effect of balneotherapy (both as a solitary approach and in the context of spa) on various diseases.	29 RCTs were evaluated; 22 investigated the use of balneotherapy in rheumatological diseases: eight osteoarthritis, six fibromyalgia, four ankylosing spondylitis, four Rheumatoid Arthritis (RA) and three RCTs in other Musculoskeletal System diseases (MSD) and chronic low back pain. 1720 patients were evaluated in these studies. Balneotherapy did result in reduced pain in patients with rheumatological diseases and chronic low back pain, in comparison to the control group in 17 of the 25 RCTs examined. This beneficial effect was measured differently and lasted for different periods of time; from ten days to one year. The available data suggest that balneotherapy may be truly associated with improvement in several rheumatological diseases.
Matz H. et. al. Balneotherapy in dermatology, <i>Dermatologic Therapy</i> , Vol. 16, 2003	<i>Evidence overview.</i> An article describing the evidence behind balneotherapy, which involves immersion of the patient in mineral water baths or pools. The major importance of balneotherapy and spa therapy both individually and as complements to other therapies lies in their potential after standard medical treatments have failed to give comfort to these patients.	<p>Bathing in water with a high salt concentration is safe, effective, and pleasant for healing and recovery. This approach needs no chemicals or potentially harmful drugs. There are almost no side effects during and after treatment, and there is a very low risk to the patient's general health and well-being.</p> <p>The major dermatologic diseases that are frequently treated by balneotherapy with a high rate of success are psoriasis and atopic dermatitis. The mechanisms behind spa therapy have not been fully elucidated. They probably incorporate chemical, thermal, mechanical, and immunomodulatory effects.</p>
Scherba G. et. al. Quantitative Assessment of the Germicidal Efficacy of Ultrasonic Energy Applied and <i>Environmental Microbiology</i> , July 1991	<i>Laboratory Study.</i> The purpose of this study was to evaluate quantitatively the degree to which microorganisms could be inactivated through the administration of ultrasonic energy into water, 39 +/- 0.3°C. Aqueous suspensions of specific bacteria, fungus, and viruses were exposed to an ultrasonic frequency of 26 kHz. The pathogens were selected based on their normal routes of infection (e.g., skin or intestinal tract) or their structural similarities to such agents, likely to be found as contaminants of whirlpools and hot tubs.	There was a significant effect on all four bacteria, with percent killed increasing with increased duration of exposure. There was a significant reduction in fungal growth compared with that in the controls, with decreased growth with increased ultrasound intensity. There was a significant reduction for feline herpesvirus with intensity, but there was no apparent effect of ultrasound on feline calicivirus. These results suggest that ultrasound in the low-kilohertz frequency range is capable to some degree of inactivating certain disease agents that may reside in water. The physical mechanism of inactivation appears to be transient cavitation.

Bathing at the bedside

A bed bath is not the most effective way of washing patients, however it may sometimes be the chosen solution. For some patients who are unable to transfer out of the bed it may be the only way to meet the hygiene needs of, for example, the unconscious patient/resident or perhaps consider a patient within end of life care.

Bed bath versus showering and bathing

Bed bathing can help to maintain the hygiene needs of residents and patients who are bedridden as a result of acute illness or chronic debilitation. However, in many cases within a general care facility, the decision to provide the patient or resident with a bed bath may be taken as a result of the caregiver perception that taking a person to the bathroom takes too long.^{8,34} It can also be the chosen because the facility does not offer the right environment or equipment to allow showering or bathing to take place. Undertaking patient hygiene in bed is strong contributor to excessive static overloading on the caregiver leading to pain and injury.^{33,34} Additional studies have shown that the

time caregiver spends in an unhealthy posture is higher in the activity of bathing at the bedside when compared to showering in a height adjustable shower chair or trolley. Knibbe et al (2016). If the decision to bathe a resident/patient within the bed is taken then consideration of the use of 'washing without water' options, that is the use of pre-heated wipes impregnated with a special cleansing and caring lotion have been shown to both improve quality of care by increasing patient satisfaction as well as increasing efficiency of work for the caregiver when compared to the traditional process of using soap, water and towels.³⁴

Using wash gloves led to a reduction in the number of repositioning activities as opposed to the traditional bed bathing method with water, soap, washcloths, and towels. The reduction of postural load for caregivers was also significant as less time was spent in ergonomically unsound postures. An increase in efficiency was found as a full body wash took an average 7.2 minutes less time than the traditional method.³⁴

A final thought....

In this clinical evidence brochure we have indicated many of the challenges facing caregivers, patient and residents in providing and receiving hygiene care. Concerns such as caregiver workplace strain and injuries, time inefficiencies, equipment and environment provision as well as the concern for the patient comfort, choice and general well being all influencing the hygiene care process. We realise that you have a choice when it comes to your provider of hygiene solutions, and also recognise that using the right equipment represents only one important part of effective hygiene care. With adequate equipment and changes to the hygiene environment, safer, calmer and more ergonomic hygiene procedures for both the resident, patient and caregiver are indeed possible.

With Arjo's many years of knowledge and experience with hygiene solutions we can also support your facility with a range of education, assessment and service solutions. For further information please visit www.arjo.com. We can help you to get the right product, to the right patient, at the right time! We welcome your questions and invite you to contact us through your local representative or via our website.

References

- 1 United Kingdom Office for National Statistics, 3rd November 2015, reference number 004835.
- 2 Freitag S, Ellegast R, Dulon M et al (2007). Quantitative measurement of stressful trunk postures in nursing professions. *Annals of Occupational Hygiene*, 51(4): 395-395.
- 3 CEN//ISO TR 12296 – 2013 Ergonomics - Manual Handling of People in the Healthcare Sector.
- 4 CDC (2011). Non-fatal bathroom injuries among persons aged > 15 years – United States, 2008. *Morbidity and Mortality weekly report*, 60(22): 729-730.
- 5 Prevention and Treatment of Pressure Ulcers: Clinical Practice Guidelines, NPUAP, EPUAP, PPPIA 2014.
- 6 Downey L, Lloyd H (2008) Bed bathing patient in hospital. *Nursing Standard* 22 (34) Pages 35-40
- 7 Brinkhoff A., Knibbe N, (2003) The ErgoStat Program. Pilot study of an ergonomic intervention to reduce static loads for caregivers, www.asse.org Professional Safety, Ergonomics, 32-39.
- 8 Knibbe JJ, Knibbe NE, Heitink DEBL (2016) Evaluating different methods of showering and washing patients: assessing ergonomic, time, and quality aspects. *American Journal of Safe Patient Handling*. Vol 6, Number 2, 49-64
- 9 Knibbe NE, Knibbe JJ, Waijjer E (2013) How smart is the Carevo? Results of a study on a new shower trolley generation. *The Ergonomics Open Journal*, 6, 1-5.
- 10 Schimmel G, Knibbe NE, (2008) The Smart Shower Chair. Final Report of the Better Care Smart Shower Chairs Project.
- 11 Hallström K. Wiese N. (2015) Improving Quality of Care and Workflow by Optimising the Working Conditions. A Case Study from a Danish Nursing Home. Proceedings 19th Triennial Congress of the IEA, Melbourne 9-14 August 2015)
- 12 Goto Y, Hayasaka S, Kurihar S et al. Physical and Mental Effects of Bathing: A Randomized Intervention Study. *Evid Based Complement Alternat Med*. 2018 (published online).
- 13 The Arjo Architect Guidebook (2019)
- 14 Freitag, S., Seddouki, R., Dulon, M., Keresten, J., Larsson, TJ., Nienhaus, A. The effect of working position on trunk posture and exertion for routine nursing tasks: an experimental study. *Ann Occup Hyg*. 2014;59(3):317-325
- 15 Branco M, Rêgo NN, Silva PH et al. Bath thermal waters in the treatment of knee osteoarthritis: a randomized controlled clinical trial. *Eur J Phys Rehabil Med*. 2016;52(4):422-30.
- 16 Kohara K, Tabara Y, Ochi M et al. Habitual hot water bathing protects cardiovascular function in middle-aged to elderly Japanese subjects. *Sci Rep*. 2018; 8: 8687.
- 17 Sakurai R1, Yasunaga M, Saito K et al. Effects of a comprehensive intervention program, including hot bathing, on physical function in community-dwelling healthy older adults: a pilot randomized controlled trial. *Aging Clin Exp Res*. 2013; 25(4): 453-61
- 18 Knibbe NE, Knibbe JJ (1996). Postural Load of nurses during bathing and showering of patients: Results of a laboratory Study. *American Society of Safety Engineers*. November. 37-39
- 19 Fujimoto S, Iwawaki Y, Takishita Y et al. Effects and safety of mechanical bathing as a complementary therapy for terminal stage cancer patients from the physiological and psychological perspective: a pilot study. *Jpn J Clin Oncol*. 2017 Nov 1;47(11):1066-1072)
- 20 Matzer F, Nagele E, Bahadori B et al (2014). Stress-relieving effects of short-term balneotherapy - a randomized controlled pilot study in healthy adults. *Forsch Komplementmed* 21(2): 105-10.)

- 21 Silva A, Queiroz SS, Andersen ML, et al (2013). Passive body heating improves sleep patterns in female patients with fibromyalgia. *Clinics (Sao Paulo)* 68(2): 135-139.
- 22 Al-Qubaeissy K, Fatoye F, Goodwin P et al (2013). The effectiveness of hydrotherapy in the management of rheumatoid arthritis: A systematic review. *Musculoskeletal Care*, 11(1): 3-18.
- 23 Staczkowski JM, Langridge J, Oatway J, Wilson C ((2018). Effect of a spa bath on patient symptoms in an acute palliative care setting: a pilot study. *Complimentary Therapies in Clinical Practice*. Accepted Manuscript
- 24 Sung HC, Chang AM. Use of preferred music to decrease agitated behaviours in older people with dementia: a review of the literature. *J Clin Nurs*. 2005 Oct;14(9):1133-40.
- 25 Sound (music): Ray KD, Fitzsimmons S. Music-assisted bathing: making shower time easier for people with dementia *J Gerontol Nurs*. 2014 Feb;40(2):9-13.
- 26 Fazaa A, Souabni L, Ben Abdelghani K et al (2014). Comparison of the clinical effectiveness of thermal cure and rehabilitation in knee osteoarthritis. A randomized therapeutic trial. *Annals of Physical and Medical Rehabilitation Medicine*, 57(9-10): 561-9
- 27 Tefner I, Kovacs C, Gaal R et al (2015). The effect of balneotherapy on chronic shoulder pain. A randomised controlled single blind follow up trial. A pilot study. *Clinical Rheumatology*, 34(6): 1097-8
- 28 Lim KO, Lee DY, Shin WS. The effects of a warm whirlpool bath on pain and stiffness of patients with chronic stroke induced knee osteoarthritis. *J Phys Ther Sci*. 2013 Jul;25(7):873-5.
- 29 Wollina U, Heinig B, Naumann G et al. Effects of low-frequency ultrasound on microcirculation in venous leg ulcers. *Indian J Dermatol*. 2011; 56(2): 174-179
- 30 Nichter LS, McDonald S, Gabriel K et al. Efficacy of debridement and primary closure of contaminated wounds: a comparison of methods. *Ann Plast Surg*. 1989; 23: 224-230
- 31 Nichter LS, Williams J. Ultrasonic wound debridement. *Journal Hand Surgery*. 1988; 13A: 142-146
- 32 Ray KD, Fitzsimmons S, Music assisted bathing, making shower time easier for people with dementia. *Journal of Gerontology Nurse* (2014) Feb 40 (2) 9-13
- 33 Myny, D., De Bacquer, D., Van Hecke, A., Beeckman, D., Verhaeghe, S., Van goubergen D. Validation of standard times and influencing factors during the development of the Workload Indicator for Nursing. *J Adv Nurs*. 2014;70 (3):674-686
- 34 Knibbe, JJ., Matz, M., & Heitink, D. (2017). Effects of Using Bathing Gloves on Healthcare Recipients and Caregivers. *Int J of SPHM*. 7(1):9-19.

Only Arjo designed parts, which are designed specifically for the purpose, should be used on the equipment and products supplied by Arjo.
As our policy is one of continuous development we reserve the right to modify designs and specifications without prior notice. © Arjo, March 2019.

At Arjo, we are committed to improving the everyday lives of people affected by reduced mobility and age-related health challenges. With products and solutions that ensure ergonomic patient handling, personal hygiene, disinfection, diagnostics, and the effective prevention of pressure ulcers and venous thromboembolism, we help professionals across care environments to continually raise the standard of safe and dignified care. Everything we do, we do with people in mind.

Arjo AB • Hans Michelsensgatan 10 • 211 20 Malmö • Sweden • +46 10 335 4500

www.arjo.com